

Contra Costa County
DECISION DOCUMENTATION for GOPHER MANAGEMENT in LANDSCAPES

Date: 5/12/16

Department: Public Works Grounds Division and Special Districts

Location: Countywide

Situation: Gophers in parks, frontage landscaping, and County landscaping

<p>What is the management goal for the sites?</p>	<p>Gopher management in the County does not seek to eradicate the animals. The management goals are to prevent gopher damage to landscaping and to building foundations or other infrastructure such as irrigation pipes and tubing, and prevent tripping hazards where children, adults, and pets play. Historically, there was such a large population of gophers in the area above Reliez Valley Rd. in the Hidden Pond Landscaping Zone that gophers were being controlled to minimize destabilization of the slope to prevent landslides.</p>
<p>Who has jurisdiction over the areas in question?</p>	<p>The County has jurisdiction over the sites; however, in Special District frontage or other landscaping, the County does not control the allocation of funds for landscape maintenance, including pest management.</p> <p>Note that Special District landscaping zones formed before 1996 do not have a built-in CPI escalator, which makes it difficult to increase the funding available for landscape maintenance. The 3 zones currently monitored for gophers are Livorna Park, Hidden Pond Landscaping Zone, and Driftwood Landscaping Zone. Hidden Pond was formed in 1990, and Driftwood was formed in 1993.</p>
<p>How often are the sites monitored?</p>	<p>This varies from site to site.</p> <p>In the course of their other work, Grounds Division staff survey for evidence of gophers. The Division also responds to complaints about gophers from County staff.</p> <p>The vertebrate pest management contractor for Special Districts regularly surveys for gophers in Livorna Park, Hidden Pond Landscaping Zone, and Driftwood Landscaping Zone and responds to complaints relayed through Special Districts staff.</p>
<p>The problem species has been identified as the following:</p>	<p>Pocket gopher, <i>Thomomys</i> sp.</p> <p>From the UC IPM Pest Notes on pocket gophers (http://www.ipm.ucdavis.edu/PMG/PESTNOTES/pn7433.html):</p> <p>“Pocket gophers are herbivorous and feed on a wide variety of vegetation but generally prefer herbaceous plants, shrubs, and trees. Gophers use their sense of smell to locate food. Most commonly they feed on roots and fleshy portions of plants they encounter while digging. However, they sometimes feed aboveground, venturing only a body length or so from their tunnel opening. Burrow openings used in this manner are called ‘feed holes.’ You can identify them by the absence of a dirt mound and by a circular band of clipped vegetation around the hole. Gophers also will pull entire plants into their tunnel from below. In snow-covered regions, gophers can feed on bark several feet up a tree by burrowing through the snow.</p> <p>“...A single gopher moving down a garden row can inflict considerable damage in a very short time. Gophers also gnaw and damage plastic water lines and lawn sprinkler systems. Their tunnels can divert and carry off irrigation water, which leads to soil erosion. Mounds on lawns interfere with mowing equipment and ruin the aesthetics of well-kept turfgrass.”</p> <p>Gophers sometimes girdle trees and shrubs and can kill trees with trunks several inches in diameter.</p> <p>Gophers also mix, aerate, and loosen soil, all of which can promote plant growth.</p>
<p>What is the tolerance level for this species?</p>	<p>One gopher burrowing in ornamental landscaping or a lawn will trigger management actions. Gophers in adjacent fields or in areas that are more wild are not managed except where gophers become numerous enough to destabilize the hillsides. Currently this applies to Hidden Pond Landscaping Zone only.</p>

Are these sensitive sites?	Are any areas part of the court-ordered injunctions? (see: https://www.epa.gov/endangered-species/interim-use-limitations-eleven-threatened-or-endangered-species-san-francisco-bay)	No for the 2 sites where rodenticide was used in the past: Hidden Pond and Driftwood.
	Are any of the sites known or potential habitats for any endangered or threatened species?	No
	Are any of the sites on or near an area where people walk or children play? Care must be taken when using gopher traps, so that neither pets nor children are likely to encounter them.	Yes
	Are any of the sites near a drinking water reservoir?	Not applicable
	Are any of the sites near a creek or flood control channel?	Not applicable
	Are any of the sites near crops?	No
	Are any of the sites near desirable trees or landscaping?	Yes
	Are any of the sites on soil that is highly permeable, sandy, or gravelly?	Not applicable
	At any of the sites, is the ground water near the surface?	Not applicable
	Are there any well heads near the sites?	Not applicable
	What factors are taken into account when determining the management technique(s) for gophers?	The proximity to foot traffic—currently traps are not used where children or other passersby might find and try to remove or tamper with the trap. Other considerations are the following: safety to the gopher manager, the environment, and non target species; endangered species considerations; the effectiveness of the method; and the cost to the Department or the Special District.
What factors contribute the cost of gopher management?	<ol style="list-style-type: none"> 1. The number of gophers at the site. 2. The number of gopher mounds at the site—each must be tamped down to determine which tunnels are active. 3. The size of the site—if a large site must be surveyed on foot, it will take longer. 4. The distance of the site from the corporation yard. 5. The skill and experience of the pest manager—someone with little experience and skill will take longer to find and trap gophers or kill them with CO₂. 6. The frequency of re-invasion—sites near open fields, vacant lots, construction sites, and wildlands will experience repeated gopher invasions. 	
Are special permits required to trap or otherwise kill gophers?	No special permits are required. Gophers are considered nongame animals by the California Department of Fish and Wildlife, which means that if a property owner finds gophers that are injuring garden or landscape plants or other property, the property owner can control the gophers at any time in any manner that is legal.	
Which cultural controls were considered?	<p>Flooding: This method is not particularly effective and would use large amounts of precious water. Most gophers survive flooding in their burrows. Some may be forced to the surface, but the pest manager would have to use something like a shovel to kill those exiting burrows.</p> <p>Planting buffers or repellent plants: A 50 ft. buffer planted in a grain, such as wheat, is mentioned in the literature, but this is not practical for the County. There is no evidence for the efficacy of planting so-called gopher repellent plants such as castor bean.</p>	

	<p>Conclusion: There are no practical or effective cultural controls for gophers in County landscaping.</p>
<p>Which physical controls were considered?</p>	<p>Trapping: Trapping is a very effective management method. There is skill and art to trapping, especially in finding the proper burrow in which to place traps; therefore, the more experienced the trapper, the more successful they are. Each management situation is unique and must be assessed at the time of inspection to determine a plan of action.</p> <p>There are a number of styles of gopher traps. The Grounds Division uses the Victor Black Box Trap. The Special District contractor uses the Gophinator trap, and the GopherHawk trap.</p> <ul style="list-style-type: none"> • The gopher manager surveys the area to determine which gopher mounds look the freshest and flags those mounds. The remaining mounds are flattened. • The following day, the manager returns to determine which mounds are actually the newest. Brand new mounds, or mounds that had been flattened and were then pushed up again, indicate the gopher is working in those areas. Otherwise the flagged mounds are still the most recent. • Working near the newest mounds, the manager uses a probe (a long pole) to find the main gopher tunnel. • A small area above the main tunnel is excavated so the traps can be inserted. Two traps are set, one in each direction back to back, so that a gopher travelling along the tunnel in either direction will encounter the business end of the trap. • The hole is covered with a board. Recommendations vary on whether or not to cover the hole, and some sources indicate that it doesn't matter, but in the County, the hole should be covered to help prevent the public from investigating the trap. The spot is marked with a small flag. <p>In an April 2013 paper in <i>Crop Protection</i>, Baldwin, et al. found that the Gophinator trap was more effective than the Macabee trap [another similar body gripping trap], probably because it was able to capture larger gophers. They also found that covering traps in late spring to early summer increased catches, but not during autumn. They recommended that if efficacy is paramount, traps should be covered from late spring to early summer, but if time is a constraining factor, traps can be left uncovered.</p> <ul style="list-style-type: none"> • Sometimes gophers are trapped immediately while the manager is still working at the site. If not, the manager returns within 24 hours to check the traps. <p>Explosive Devices: The Rodenator injects a combination of 3% propane and 97% oxygen into a burrow and ignites these gases. The resulting explosion collapses the tunnel and creates a shockwave that kills gophers in the burrow. Around 2013, the Grounds Division conducted a trial of the Rodenator outside the Public Works Administration building on Glacier Drive in Martinez. Gophers were burrowing close to the building, and it was feared that they might undermine the foundation. The device worked well and no gophers have been seen in that area since. There are, however, some problems with this device. All the windows on the treatment side of the building had to be protected with sheets of plywood, and the explosions rattled the windows and the occupants of the building. The reports from the explosions, which sound like gunshots, precipitated calls to the police, even though the surrounding neighbors had been notified. The Division has not pursued this strategy because of this last issue. There is also a fire risk with this method.</p> <p>Exclusion with wire mesh: Three-foot high ½" wire mesh buried 2 feet below ground and encircling a plant can exclude gophers temporarily. These wire cages are only effective in protecting a small area and are very expensive to make and install.</p> <p>Conclusion: Trapping is the most effective and practical physical control for gophers in County landscaping. All gopher problems are currently managed with trapping.</p>
<p>Which biological controls were considered?</p>	<p>Great blue herons, coyotes, domestic dogs and cats, foxes, and bobcats capture gophers at their burrow entrances; badgers, long-tailed weasels, skunks, rattlesnakes, and gopher snakes corner gophers in their burrows. Owls and hawks capture gophers above ground.</p> <p>Predators can prune a population, but none of these predators can control gophers to the extent that is necessary in County landscaping. Owl boxes could attract more owls to certain areas of the County. More owls could mean somewhat fewer gophers in open fields.</p> <p>Conclusion: Biological controls alone for gophers have not been shown to reliably reduce populations to the level that will prevent damage to plants and infrastructure.</p>

<p>Which chemical controls were considered?</p> <p>For more information on pesticides listed here visit the National Pesticide Information Center (NPIC). This a joint project of Oregon State University and the US EPA.</p> <p>http://npic.orst.edu/</p> <p>You can communicate with an actual person at 1.800.858.7378 or npic@ace.orst.edu</p> <p>They are open from 8:00AM to 12:00PM Pacific Time, Mon-Fri</p>	<p>The risk to predatory animals must be considered before any rodenticides are used for gopher management.</p> <p>Fumigants</p> <p>Extension and university literature recommend against using fumigants for gophers because the animals can quickly backfill a tunnel when they perceive a threat, which prevents the gas from reaching them. Injecting gas far enough into their extensive burrow system is difficult, and since their tunnels are close to the surface, gas can leak out and never reach a concentration high enough to kill.</p> <p>CO₂ Injection</p> <ul style="list-style-type: none"> • The Grounds Division has purchased a device called the Eliminator which injects carbon dioxide into the burrow system. So far the gopher manager has had good luck with this device. Perhaps this is more effective since the CO₂ initially sinks to the floor of the burrow. • This device can be used where foot traffic prohibits the use of traps. • The same preliminary procedures are employed for this device as for trapping (see above). • Before deploying the device in the burrow, any openings should be closed and remaining mounds should be flattened to help keep the gas inside the burrow. • When the trigger on the device is pulled, there should be no hissing sounds. • The area should be monitored the day after the treatment to determine the degree of success. <p><i>A note on "signal words," below: these designations from the USEPA pertain to the acute toxicity of a pesticide.</i></p> <p>Aluminum Phosphide Signal Word: DANGER</p> <ul style="list-style-type: none"> • Fumigation with aluminum phosphide <u>is</u> effective for gophers, although it is a restricted use material that requires a permit from the County Department of Agriculture. Aluminum Phosphide is not used in the County for gophers. <p>Baiting</p> <p>Diphacinone (005%) Multiple Dose Bait Blocks (Eaton's Answer®) Signal Word: CAUTION. Baiting is no longer used for gophers in Contra Costa County.</p> <p>Conclusion: CO₂ injection has worked well for the Grounds Division, but lack of staff has curtailed its use. For large areas with many ground squirrels, it could be used again.</p> <p>Baiting is not being used.</p>
<p>Recommendations from the IPM Advisory Committee</p>	<p>On-going monitoring should be used to adjust control activities to a level appropriate to the population of gophers. Trapping and CO₂ injection are the preferred control methods when sufficient funding is available.</p> <p>Consider expanding trapping into areas where children or other passersby have access after investigating techniques used in school IPM programs or other programs where trapping is conducted in sensitive sites.</p>
<p>References</p>	<p>Baldwin, R.A., D.B. Marcum, S.B. Orloff, S.J. Vasquez, C.A. Wilen, and R.. Engeman (2013). The influence of trap type and cover status on capture rates of pocket gophers in California, <i>Crop Protection</i>, 46: 7-12.</p>